



## Potentials of hydrogels in rainfed soil to conserve soil moisture and fertility to maximize the wheat yield

Aftab Ahmad Shaikh<sup>1</sup>, Obaid ur Rehman<sup>2</sup>, Muhammad Rashid<sup>2\*</sup>, Sarosh Alvi<sup>2</sup>, Abad Raza<sup>2</sup>, Abdul-Waheed<sup>2</sup>, Muhammad Irfan<sup>3</sup>, Shahid Saleem<sup>2</sup>, Muhammad Muddassir Iqbal<sup>3</sup> and Ghulam Mujtaba<sup>3</sup>

<sup>1</sup>Pesticide Residue Laboratory, Kala Shah Kaku

<sup>2</sup>Soil and Water Testing Laboratory, Rawalpindi

<sup>3</sup>PMAS-Arid Agriculture University, Rawalpindi

[Received: August 21, 2020 Accepted: October 01, 2020 Published Online: November 28, 2020]

### Abstract

The availability of limited irrigation water in rainfed regions is the most challenging constraint to harvest the maximum agronomic benefits. The adaptation of smart water-saving techniques is desperate to combat soil moisture stress and soil quality. The application of hydrogels to conserve soil moisture and plant nutrients to satisfy the crop needs is considered one of the most efficient strategies. Keeping in view the peculiar benefits of hydrogels, an experiment was planned and executed in rainfed areas to quantify the potential benefits of hydrogels for the conservation of soil moisture and restoration of soil fertility under wheat production compared with various conventional soil amendments. The treatments were: T<sub>1</sub> (Control), T<sub>2</sub> Farmyard manure @ 2500 kg ha<sup>-1</sup>, T<sub>3</sub> Compost @ 750 kg ha<sup>-1</sup>, T<sub>4</sub> Gypsum @ 1000 kg ha<sup>-1</sup>, T<sub>5</sub> Hydrogel @ 15 kg ha<sup>-1</sup>. The findings of this experiment revealed that soil moisture retention increased from 27 to 73% compared with control (untreated soil) and the application of soil amendments enhanced the soil NO<sub>3</sub>-N (6 to 24 mg kg<sup>-1</sup>), P<sub>2</sub>O<sub>5</sub> (4.75 to 11 mg kg<sup>-1</sup>), and K<sub>2</sub>O (93.50 to 115.25 mg kg<sup>-1</sup>), respectively, by farmyard manure, compost, gypsum and hydrogel under specified doses. The wheat grain yield increased from 2165 to 3760 kg ha<sup>-1</sup>. This study has provided preliminary results regarding the use of hydrogels in a field experiment in rainfed areas of Pakistan. Further studies are required to investigate the performance of various hydrogels under variable environments before recommending to the farmers.

**Keywords:** Soil amendments, water use efficiency, fertilizer use efficiency, water holding capacity; rainfed areas

### Introduction

The agricultural production in rainfed areas is confronted with many challenges including uncertain frequency and the distribution of rainfall. The success of crops largely depends upon the water supply through rains otherwise soil moisture shortage will result in drought stress (Abdelfattah, 2013). Various management strategies including the conventional soil amendments have been tested but could not offer a sustainable solution to defer moisture stress and fertility depletion. The most sustainable solution of all these issues is the prolonged retention of rainwater through appropriate water conservation techniques to ensure the crop survival (Bhardwaj *et al.*, 2007) since the limited water supply in rainfed areas is the main constraint to achieve the successful crop (Han *et al.*, 2018). The moisture deficiency accelerates the decomposition of soil organic matter causing poor water holding capacity and nutrient use efficiency (Mandal *et al.*, 2011). The traditional

approaches including the addition of a large quantity of organic biomass are practiced but the adoption of such practices on large scale is not viable (Ankenbauer and Loheide, 2017; Minasny and McBratney, 2018).

Several soil amendments such as gypsum, organic materials, and hydrogels, etc. have been evaluated by the researchers to conserve the profile soil moisture and plant nutrients but most of these could not gain widespread acceptance due to their short-term benefits and high cost. However, many studies have proved that the addition of hydrogels as a soil amendment is effective in enhancing soil water-holding-capacity by minimizing the excessive water drainage and evaporation (Cannazza *et al.*, 2014; Guilherme *et al.*, 2015). The hydrogels are channels of polymers having the ability to hold large quantities of water (Azevedo *et al.*, 2016). The incorporation of a precise number of hydrogels in the soil can improve water holding capacity; minimize the evapotranspiration and, mitigate the drought stress symptoms (Li *et al.*, 2018). The enhanced availability

\*Email: abuxenab@gmail.com

of water may assist to defer the water stress condition during long periods of drought. It is assumed that water release through the network of hydrogel will develop additional space in the soil which will facilitate the root growth and water penetration for long-term storage. Additionally, this stored water will restrict nutrient losses through leaching and percolation, consequently, extending the supply of water and nutrients for an extended period (Karimi and Naderi, 2007; Bagheri and Afrasiab, 2013; Pattanaik *et al.*, 2015). Furthermore, the addition of hydrogels may enhance the fertilizer use efficiency in soil by extending the nutrient release period (Zhang *et al.*, 2006).

Hydrogels can be categorized into three groups of a polymer including i) Starch polyacrylonitrile graft polymers (starch copolymers), ii) Vinyl alcohol-acrylic acid copolymers (polyvinyl alcohols), and iii) Acrylamide sodium acrylate copolymers-cross-linked polyacrylamides. Hydrogel-forming natural polymers include proteins such as collagen and gelatin and polysaccharides such as starch, alginate, and agarose. Synthetic polymers that form hydrogels are traditionally prepared by using chemical polymerization methods (Ahmed, 2015). The hydrogel selected for this study is known as polyacrylamide (Qemisoyl) which can absorb and retain a large amount of water along with nutrients for the uptake of crops. One gram of this hydrogel can absorb up to 500 mL of water and is recommended to be applied once and remain functional for 4-7 years in the soil (Monning, 2005).

The topography of the area under this study is sloppy which limits the water infiltration and storage due to excessive runoff. Several scientists have revealed that the slope gradient can increase soil erosion due to poor infiltration rate and enhance the surface run off (Neethu *et al.*, 2018). Pakistan is an agricultural country with having diverse climate and about two-thirds of its lands are categorized as arid facing the high temperature and low rainfall (Chaudhry and Rasul, 2004). Wheat is the most important crop being the staple food, but it faces at least two dry spells because of uncertain rainfall under the changing climate scenario. It is predicted that Pakistan will receive lesser winter rains in the coming times which may delay the sowing of wheat under rainfed areas of Pakistan. The sowing of wheat is carried out in October to November, but the weather forecast has predicted a reduction in winter rains (Hanif and Ali, 2014). An ample supply of irrigation water for crops during the growth and development of crops is compulsory for the successful crop to stand to ensure food security in the region (Iqtidar *et al.*, 2006). So, the innovative approaches to enhance water retention especially for wheat production under

rainfed conditions are needed desperately to achieve the desired yield goals. Many studies across the world have proved the potential benefits of hydrogels as a soil amendment but there is no information available in rainfed areas of Pakistan regarding the use of hydrogels especially for field crops. So, there is a need for extensive research to investigate the potential benefits of hydrogels and the standardization application rate.

Therefore, considering the benefits of hydrogels and other soil conditioners and peculiar problems of rainfed areas, a field study was undertaken to compare the benefits of these soil conditioners by evaluating their impact on soil chemical properties and wheat yield.

## Materials and Methods

### Location

The location of the experiment selected in Fateh Jang (Attock) was situated on latitude; 33.55° N, and longitude; 72.58° E. The majority of soils of the area are prone to soil erosion, gully development, and soil moisture shortage. The climate is semi-arid to sub-tropical receiving 584 mm mean annual rainfall, out of which only 30% rainfall was received during the wheat growing season. The soils are deep having good drainage (Reconnaissance Soil Survey, Campbellpur, 1970). Soil properties at the start of the experiment are given in Table-1. Before the execution of the experiment, a composite soil sample representing the characteristics of the experimental site was collected, air-dried, and separated by 2 mm sieve. The pH and EC of soil were determined by McLean (1982), and Richards (1954), Soil organic matter by Walkley (1947), K<sub>2</sub>O by Rhodes, 1982, P<sub>2</sub>O<sub>5</sub> by Watanabe and Olsen (1965), and soil particle size by Gee and Bauder (1986).

**Table-1: Soil properties at the start of experiment**

Soil property	Value	Unit
pH	8.12	-
EC <sub>e</sub>	0.76	dS m <sup>-1</sup>
Bulk density	1.59	g cm <sup>-3</sup>
O.M	0.87	%
P	3.8	mg kg <sup>-1</sup>
K	93	mg kg <sup>-1</sup>
Soil texture	Sandy loam	-

### Experiment

The study was carried out during the autumn season, 2011-12 to 2014-15. All the treatments were directly incorporated in the soil before sowing of wheat during the 1st week of November. The experiment was comprised of five treatments: T<sub>1</sub> (Control), T<sub>2</sub> FYM @ 2500 kg ha<sup>-1</sup>, T<sub>3</sub>



Compost @ 750 kg ha<sup>-1</sup>, T<sub>4</sub> Gypsum @ 1000 kg ha<sup>-1</sup>, and T<sub>5</sub> Hydrogel @ 15 kg ha<sup>-1</sup>. The experiment was laid in an RCBD arrangement with 3 replications. During the first year of study, half rate of recommended NPK (120, 80, 60 kg ha<sup>-1</sup>) was applied in hydrogel plots and full recommended dose of NPK in gypsum plots but in the following years, full doses of recommended NPK were applied before sowing of wheat in the form of urea, diammonium phosphate, and sulfate of potash. However, the amendments (gypsum and hydrogels) were applied once at the start of the study considering their long-term efficiency in soil. Wheat variety *BARS-2010* was sown every year. All the recommended agronomic and cultural practices were carried out when and where required. The experiment was conducted in rainfed conditions and no supplemental irrigation was applied except natural precipitation. The plots were harvested manually during the 1<sup>st</sup> week of May and 1 m<sup>2</sup> wheat crop sample was collected from every treatment to determine grain yield. A composite soil sample consisting of three sub-samples was also collected from each treatment for the determination of physicochemical properties. The data was analyzed statistically (using statistics 8.1) by LSD test at a 5% probability level (Steel *et al.*, 1997).

### Soil moisture retention test

The soil was collected from 0-20 cm soil layer and dried in air. The soil was ground and passed through a 2 mm sieve and 1 kg loose soil was filled in polyvinyl chloride columns having 30 cm length and 5 cm diameter. The columns were sealed from the bottom with nylon mesh cloth and placed on sealed plastic lids having small holes to drain excess water. The columns were tapped 20 times to maintain bulk density near the field conditions. All the treatments were mixed in a 5 cm top layer following the rates described in the previous section. Then, columns were saturated with distilled water from the top. Soil moisture was monitored periodically by weighing the columns at an interval of 3 days to 30 days.

## Results and Discussion

### Grain yield


The impact of soil amendments on wheat grain yield is shown in Figure 1. It was observed that all soil amendments enhanced the wheat grain yield significantly compared to control. However, when we compared the soil amendments with each other, it was noticed that the mean maximum grain yield (3760 kg ha<sup>-1</sup>) was recorded in T<sub>2</sub> while the mean minimum grain yield (2165 kg ha<sup>-1</sup>) was recorded in T<sub>1</sub>. It was estimated that overall, 42.42, 31.51, 33.91, and 34.13% increase in wheat grain yield was recorded by hydrogel, FYM, compost, and gypsum,

respectively, compared with control. Additionally, it was found that hydrogel proved the best among all soil amendments by extending an increase of 15.93, 12.87, and 12.57% in wheat grain yield compared with FYM, compost, and gypsum, respectively. Probably, the increase in wheat grain production was attributed to an increase in better supply of essential nutrients after the application of soil amendments. The best results obtained due to hydrogel application confirms the findings of Eiasu *et al.* (2007) who found that application of hydrogels with fertilizers, improved the crop yield. El-Hardy *et al.* (2009) also reported a significant increase in wheat yield and investigated that the soil moisture retention extended the supply of nutrients by the hydrogels under water stress conditions which could have resulted in better crop yield. It was also found from the data of this study that the performance of other soil conditioners i.e. FYM and compost was declined with the passage of time could be due to the rapid decomposition of soil organic matter supplied by these amendments. While the performance of gypsum was ranked second after the hydrogels could be due to enhanced soil water infiltration and improved soil structure resulted in better nutrient use efficiency (Rehman *et al.*, 2013). The application of hydrogels increased the wheat grain yield, might be a result of better assimilation of carbohydrates in the ear. The ear formation is also directly related to the number of productive tillers and conducive condition for the formation of more number of productive tillers such as the increase in CO<sub>2</sub> assimilation rate, delay in senescence of flag leaf, and effective translocation of dry matter from source to sink which together resulted in the production of a higher number of spikes with longer spike length (Tyagi *et al.*, 2015).


### Soil fertility attributes

The data of NO<sub>3</sub>-N concentration in soil is presented in Figure 2. The data obtained in this experiment has shown that the mean maximum soil NO<sub>3</sub>-N contents (24.25 mg kg<sup>-1</sup>) were measured in T<sub>2</sub> followed by 15.75 mg kg<sup>-1</sup> in T<sub>4</sub> while the mean minimum soil NO<sub>3</sub>-N concentration (6.00 mg kg<sup>-1</sup>) was noticed in T<sub>1</sub>. The data showed that there was an increase of 75.25, 59.32, 62.02, and 46.66% after the application of hydrogel, gypsum, compost, and farmyard manure, respectively, compared to control. The best performance of hydrogel treatment could be due to the fact that hydrogel might have entrapped the soil N from the soil solution which has minimized its losses in the form of leaching and denitrification. Hydrogel application reduced the nutrient loss and enhanced the nutrient use efficiency by reducing the need for fertilization (Ni *et al.*, 2009).






**Figure 1: Impact of soil amendments on wheat grain yield**



**Figure 2: Impact of soil amendments on soil NO<sub>3</sub>-N concentration**


The results of this study also revealed that the incorporation of soil conditioners in soil significantly enhanced the  $P_2O_5$  contents in all treatments compared with control however, the hydrogel performed the best and showed mean maximum  $P_2O_5$  contents ( $11.00 \text{ mg kg}^{-1}$ ) followed by gypsum whereas, the mean minimum soil  $P_2O_5$  contents ( $4.75 \text{ mg kg}^{-1}$ ) were recorded in  $T_1$  (control). It was concluded that soil conditioners including hydrogel, FYM, compost, and gypsum have increased the  $P_2O_5$  contents by 56.81, 34.48, 48.64, and 50.00%, respectively compared to control. The benefits of hydrogel addition for soil  $P_2O_5$  retention in rainfed Pakistani soils could be attributed to a reduction in the runoff by sediment conservation since the loss of P through sediment movement is one of the major factors of low P use efficiency in these soils along with fixation of P by calcium at exchange sites. The guarantee of a sustainable supply of available soil moisture for an extended period helps in the development of soil aggregates which ensures better soil quality and P bioavailability in calcareous soils. This may also reduce the runoff by enhancing the water infiltration rate. Previous studies indicated that the addition of hydrogels @  $10 \text{ mg L}^{-1}$  with irrigation water reduced the loss of total P by up to 92% (Lentz *et al.*, 2001), while under sprinkler irrigation

polyacrylamide (PAM) reduced the loss of P applied @  $2\text{--}4 \text{ kg ha}^{-1}$  (Bjorneberg *et al.*, 2000). It was also investigated that lower rates of PAM were more effective in irrigated agriculture, could be due to direct application with irrigation water. But, in rainfed agriculture, higher application rates of PAM ( $>20 \text{ kg ha}^{-1}$ ) are needed to conserve soil moisture for a longer period since using hydrogel as a soil conditioner increases the CEC, specific surface area, and available nutrient status (Rajakumar *et al.*, 2016).



**Figure 3: Impact of soil amendments on soil  $P_2O_5$**

In this experiment, the researchers found that the incorporation of soil conditioners in soil significantly improved the status of soil  $K_2O$  concentration in all treatments compared with control. However, the addition of hydrogels has revealed the best response and showed the mean maximum  $K_2O$  concentration ( $115.25 \text{ mg kg}^{-1}$ ) following the compost ( $101.25 \text{ mg kg}^{-1}$ ) whereas, the mean minimum soil  $K_2O$  contents ( $93.50 \text{ mg kg}^{-1}$ ) were measured in  $T_1$  (control). It was concluded that soil conditioners including hydrogel, FYM, compost, and gypsum have increased the  $K_2O$  contents by 18.87, 5.79, 7.88, and 6.03%, respectively, compared with control. The literature review pointed out that the application of gel-forming polymers enhanced the water holding and nutrient supplying capacity of soil to improve growth and dry matter production of crops (El-Hady *et al.*, 1981).




**Figure 4: Impact of soil amendments on soil  $K_2O$**



## Dynamics of soil water holding capacity

The experimental results presented in Figure 5, revealed that all soil conditioner enhanced the soil water holding capacity significantly. The maximum water holding capacity (73%) was recorded with T<sub>2</sub> under hydrogel application followed by 48% with T<sub>4</sub> where compost was applied. The accumulative minimum soil water holding capacity (27%) was measured with T<sub>1</sub> (control). However, the treatments including hydrogel, FYM, compost, and gypsum have increased the water holding capacity by 63.01, 37.20, 43.75, and 35.71%, respectively, compared to control. The comparison among soil amendments showed that the hydrogel has enhanced soil water holding capacity by 41.09, 34.24, and 42.46%, respectively, compared to FYM, compost, and gypsum, respectively.



**Figure 5: Impact of soil amendments on soil water retention**

After the application of hydrogels, the water is absorbed by osmosis and interacts with hydrogen atoms as positive ions leaving negative ions along the length of the polymer chain. This yielded several negative charges in its length which repel each other to unwind the polymer chain. The water molecules bind them with hydrogen bonding (Vicky, 2007). The application of hydrogels may serve as a slow-release water reservoir in soil that may retain soil water for extended time as these are capable of absorbing water several hundred times of their weight (Monnig *et al.*, 2005).

## Conclusion

The data obtained in this experiment showed that the application of soil conditioners including hydrogels possesses better water-retaining capacity. The addition of the hydrogel has improved water retention (73%) and enhanced wheat yield (42%). Additionally, soil conditioners have enhanced soil fertility. Therefore, it can be predicted that the application of hydrogel could be successful in maintaining stable plant growth under water stress

conditions. But the variability in the performance of hydrogels warrants further investigations for the recommendations of the specific hydrogel according to soil types.

## References

- Abdelfattah, M.A. 2013. Pedogenesis, land management and soil classification in hyper-arid environments: Results and implications from a case study in the United Arab Emirates. *Soil Use and Management* 29: 279-294.
- Ahmed, E.M. 2015. Hydrogel: Preparation, characterization, and applications: A review. *Journal of Advanced Research* 6: 105-121.
- Ankenbauer, K.J. and S.P. Loheide, 2017. The effects of soil organic matter on soil water retention and plant water use in a meadow of the Sierra Nevada, CA. *Hydrological Processes* 31(4): 891-901.
- Azevedo, G., G.B.de-Azevedo, A.M. de-Souza, C.L. Mews and J.R.L. de-Sousa. 2016. Effect of hydrogel doses in the quality of *Corymbia citriodora* Hill & Johnson seedlings. *Nativa* 4: 244-248.
- Bagheri, H. and P. Afrasiab. 2013. The effects of super-absorbent, vermicompost and different levels of irrigation water salinity on soil saturated hydraulic conductivity and porosity and bulk density. *International Research Journal of Applied and Basic Sciences* 4(8): 2381-2388.
- Bhardwaj, A.K., I. Shainberg, D. Goldstein, D. Warrington and G.J. Levy. 2007. Water retention and hydraulic conductivity of cross-linked polyacrylamides in sandy soils. *Soil Science Society of America Journal* 71 (2): 406-412.
- Bjorneberg, D.L., J.K. Aase and D.T. Westermann. 2000. Controlling sprinkler irrigation runoff, erosion, and phosphorus loss with straw and polyacrylamide. *Transacts of ASAE* 43(6): 1545-1551.
- Cannazza, G., A. Cataldo, E. De-Benedetto, C. Demitri, M. Madaghiele and A. Sannino. 2014. Experimental assessment of the use of a novel superabsorbent polymer (SAP) for the optimization of water consumption in agricultural irrigation process. *Water* 6(7): 2056-2069.
- Chaudhry, Q.Z. and G. Rasul. 2004. Agro-climatic Classification of Pakistan, Science Vision (Vol.9, No.1-2 & 3-4), (July-Dec 2003 & Jan-Jun 2004), Pakistan Meteorological Department, H-8, Islamabad. p.59.
- Eiasu, B.K., P. Soundy and P.S. Hammes. 2007. Response of potato (*Solanum tuberosum*) tuber yield components to gel-polymer soil conditioners and irrigation regimes. *New Zealand Journal of Crop Horticultural Sciences* 35(1): 25-31.



- El-Hady, O.A., M.Y. Tayel and A.A. Lotfy. 1981. Super gel as a soil conditioner, II-Its effects on plant growth, enzyme activity, water use efficiency and nutrient uptake. *Acta Horticulturae* 119:257-265.
- El-Hardy, O.A., A.A. El-Kader and A.M. Shafi. 2009. Physio-bio-chemical properties of the sandy soil conditioned with acrylamide hydrogels after cucumber plantation. *Australian Journal of Basic & Applied Sciences* 3: 3145-51.
- Gee, G. and W. Bauder. 1986. Particle size analysis. p.383-409. In: *Methods of Soil Analysis, Part I*, 2nd Ed. A. Klute (ed.). Aron. Monogr. 9. ISSSA, Madison, WI.
- Guilherme, M.R., F.A. Aouada, A.R. Fajardo, A.F. Martins, A.T. Paulino and M.F.T. Davi. 2015. Superabsorbent hydrogels based on polysaccharides for application in agriculture as soil conditioner and nutrient carrier: A review. *European Polymer Journal* 72: 365-385.
- Han, B., S.G. Benner and A.N. Flores. 2018. Evaluating impacts of climate change on future water scarcity in an intensively managed semi-arid region using a coupled model of biophysical processes and water rights. *Hydrology and Earth System Sciences Discussions* 1-53.
- Hanif, M. and J. Ali. 2014. Climate scenarios 2011-2040. Districts Haripur, Sawabi, Attock and Chakwal Pakistan. Publication, Inter-cooperation Pakistan. p. 27.
- Huang, Z.B., G.Z. Zhang, Y.Y. Li, M.D. Hao, B.H. Meni and D.L. Chen. 2002. Characteristics of Aquasorb and its application in crop production. *Transactions of Chinese Society Agricultural Engineering* 18: 22-26.
- Iqtidar, H., K.M. Ayaz and K.E. Ahmad. 2006. Bread wheat varieties as influenced by different nitrogen levels. *Journal of Zhejiang University of Sciences* 7:70-78.
- Karimi, A. and M. Naderi. 2007. Yield and water use efficiency of forage corn as influenced by superabsorbent polymer application in soils with different textures. *Agricultural Research* 7(3): 187-198. (In Persian).
- Lentz, R.D., R.E. Sojka, C.W. Robbins, D.C. Kincaid and D.T. Westermann. 2001. Polyacrylamide for surface irrigation to increase nutrient use efficiency and protect water quality. *Communication in Soil Science and Plant Analysis* 32(7-8): 1203-1220.
- Li, J.X.M., G. Sa, D. Zhou, X. Zheng, X. Zhou, C. Lu, S. Lin, R. Zhao and S. Chen. 2018. Natural and synthetic hydrophilic polymers enhance salt and drought tolerance of *Metasequoia glyptostroboides* Hu and WC Cheng seedlings. *Forests* 9: 643.
- Mandal, U.K., R. Kausalya, K.L. Sharma, B. Satyam, K. Venkanna, M.U. Bhanu, M. Mandal, R. N. Masane, B. Narsimlu, K.V. Rao, G.R.S.C. Korwar and B. Venkateswarlu. 2011. Assessing soil quality under different land-use systems in a watershed of semi-arid tropics in India using GIS. *Soil Science Society of America Journal* 75(3): 1144-1160.
- McLean, E.O. 1982. Soil pH and lime requirement. p.199-224. In: *Methods of Soil Analysis, part 2: Chemical and Microbiological Properties*. A.L. Page (ed.). American Society Agronomy, Madison, WI, USA.
- Minasny, B. and A.B. McBratney. 2018. Limited effect of organic matter on soil available water capacity. *European Journal of Soil Science* 69 (1): 39-47.
- Monnig, S. 2005. Water saturated super-absorbent polymers used in high strength concrete. *Otto- Graf- Journal* 16:193-202.
- Neethu, T.M., P.K. Dubey and A.R. Kaswala. 2018. Prospects and applications of hydrogel technology in agriculture. *International Journal of Current Microbiology & Applied Sciences* 7(5): 3155-3162.
- Ni, B., M. Liu and S. Lü. 2009. Multifunctional slow-release urea fertilizer from ethylcellulose and superabsorbent coated formulations. *Chemical Engineering Journal* 155(3):892-898.
- Pattanaaik, S.K., L. Wangchu, B. Singh, B.N. Hazarika, S.M. Singh and A.K. Pandey. 2015. Effect of hydrogel on water and nutrient management of Citrus reticulata. *Research on Crops* 16 (1): 98-103.
- Rajakumar, R. and S.J. Sankar. 2016. Hydrogel: Novel soil conditioner and safer delivery vehicle for fertilizers and agrochemicals-A review. *International Journal of Applied and Pure Science and Agriculture* 2 (9): 163-172.
- Reconnaissance Soil Survey Campbellpur. 1970. Soil Survey of Pakistan.
- Rehman, O., M. Rashid, S. Alvi, R. Kausar, R. Khalid and T. Iqbal. 2013. Prospects of using gypsum to conserve water and improve wheat yield in rainfed aridisols. *Pakistan Journal of Scientific and Industrial Research Series B: Biological Sciences* 56(1):11-17.
- Rhoades, J.D. 1982. Cation exchange capacity. P.149-158 In: *Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties*. A.L. Page (ed.). American Society of Agronomy, Madison, Wisconsin, USA.
- Richards, L.A. 1954. Diagnosis and improvement of saline and alkali soils. Handbook 60, United States Department of Agriculture, Washington 25, D.C.
- Steel, R.G.D., J.H. Torrie and M.A. Boston. 1997. Principles and Procedures of Statistics: A Biometrical Approach. McGraw Hill Book Company Inc. New York. p. 633.
- Tyagi, V., R.K. Singh and M. Nagargade. 2015. Effect of hydrogel, NPK and irrigation levels on yield, nutrient



- uptake and water use efficiency of wheat (*Triticum aestivum* L.). *Research on Crops* 16 (4): 653-656.
- Vicky, W. 2007. Hydrogels: Water-absorbing polymers. *Catalyst* 18(1): 18-21.
- Walkley, A. 1947. A critical examination of a rapid method for determining organic carbon in soils – Effect of variations in digestion conditions and of inorganic soil constituents. *Soil Science* 63(4): 251-264.
- Watanabe, F.S. and S.O. Olsen. 1965. Method for determination of phosphorus in water and  $\text{NaHCO}_3$  extracts from soil. *Soil Science Society of America Proceeding* 29: 677–678.
- Zhang, J., H. Chen, P. Li and A. Wang. 2006. Study on superabsorbent composite, 14. Preparation of poly(acrylic acid)/organo-attapulgit composite hydrogels and swelling behaviors in aqueous electrolyte solution. *Macromolecular Material Engineering* 291: 1529-1538.
- Zaker, M. and H. Emami. 2019. Effect of potassium to bivalent cations ratio in irrigation water on some physical and hydraulic properties of sandy loam soil. *Soil and Environment* 38(1): 66-74.
- Zhang, G.S., K.Y. Chan, A. Oates, D.P. Heenan and G.B. Huang. 2007. Relationship between soil structure and runoff/soil loss after 24 years of conservation tillage. *Soil and Tillage Research* 92: 122-128.

