

Use of Different Extractants for Determination of Cu Status of Bajada & terrace Plain Soils of Turkey

MUHAMMAD RAMZAN CHAUDHRY**

ABSTRACT

Twenty four soil samples (0-20cm) were collected from Bajada and Terrace plain soils. Oats were grown in pots using these soils for 5 weeks. The yield and Cu contents in the plants were, on an average, higher in Bajada plains. There was no significant correlation between dry matter yield and Cu contents of soil. The maximum amount of Cu in the soils was extracted with 0.01M EDTA + 1M(NH₄)₂CO₃ followed by 3% HNO₃ and 1NNH₄OAc.

Significant positive correlation ($r=0.586^*$) was found between Cu contents of oat plants and that of 1NNH₄OAc extractable Cu of soils of Bajada plains. No significant correlation was found between Cu contents of the oat and that of soil extracted with other extractants. Yield was not significantly correlated with Cu contents of the oat.

Significant positive correlation was found between 1NNH₄OAc-extractable Cu and soil clay ($r=0.721^*$) and CaCO₃ ($r=0.682^*$) and negative correlation ($r= -0.698^*$) between CaCO₃ and HNO₃ extractable Cu of soils of Bajada plains. In Terrace plains negative correlation was found between pH ($r= -0.776^{**}$), O.M. ($r= -0.667^*$), and NH₄OAc-extractable Cu and between pH ($r= -0.667^{**}$), CaCO₃ ($r= -0.765^{**}$) and EDTA and HNO₃ extractable Cu respectively. 1NNH₄OAc seems to be more suited extractant from these extractants.